

Neutrino Masses in the Supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ Model with right-handed neutrinos

P. V. Dong^a, D. T. Huong^a, M. C. Rodriguez^b and H. N. Long^a

^a *Institute of Physics, VAST, P. O. Box 429, Bo Ho, Hanoi 10000, Vietnam*

^b *Fundação Universidade Federal do Rio Grande, Departamento de Física,
 Av. Itália, km 8, Campus Carreiros, 96201-900 Rio Grande, RS, Brazil*

Abstract

The R-symmetry formalism is applied for the supersymmetric $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ (3-3-1) model with right-handed neutrinos. For this kind of models, we study generalization of the MSSM relation among R-parity, spin and matter-parity. Discrete symmetries for the proton stable in this model are imposed, and we show that in such a case it is able to give leptons masses at only the tree level contributions required. A simple mechanism for the mass generation of the neutrinos is explored. We show that at the low-energy effective theory, neutrino spectrum contains three Dirac fermions, one massless and two degenerate in mass. At the energy-level where the mixing among them with neutralinos turned on, neutrinos obtain Majorana masses and correct the low-energy effective result which naturally gives rise to an inverted hierarchy mass pattern. This mass spectrum can fit the current data with minor fine-tuning. Consistent values for masses of the charged leptons are also given. In this model, the MSSM neutralinos and charginos can be explicitly identified in terms of the new constraints on masses which is not as in a supersymmetric version of the minimal 3-3-1 model.

PACS numbers: 11.30.Er, 14.60.Pq, 14.60.-z, 12.60.Jv

1 Introduction

Although the Standard Model (SM) gives very good results in explaining the observed properties of the charged fermions, it is unlikely to be the ultimate theory. It maintains the masslessness of the neutrinos to all orders in perturbation theory, and even after non-perturbative effects are included. The recent groundbreaking discovery of nonzero neutrino masses and oscillations [1] has put massive neutrinos as one of evidences on physics beyond the SM.

The Super-Kamiokande experiments on the atmospheric neutrino oscillations have indicated to the difference of the squared masses and the mixing angle with fair accuracy [2, 3]

$$\Delta m_{\text{atm}}^2 = 1.3 \div 3.0 \times 10^{-3} \text{eV}^2, \quad (1.1)$$

$$\sin^2 2\theta_{\text{atm}} > 0.9. \quad (1.2)$$

While, those from the combined fit of the solar and reactor neutrino data point to

$$\Delta m_{\odot}^2 = 8.0_{-0.4}^{+0.6} \times 10^{-5} \text{ eV}^2, \quad (1.3)$$

$$\tan^2 \theta_{\odot} = 0.45_{-0.07}^{+0.09}. \quad (1.4)$$

Since the data provide only the information about the differences in m_{ν}^2 , the neutrino mass pattern can be either almost degenerate or hierarchical. Among the hierarchical possibilities, there are two types of normal and inverted hierarchies. In the literature, most of the cases explore normal hierarchical one in each. In this paper, we will mention on a supersymmetric model which naturally gives rise to three pseudo-Dirac neutrinos with an inverted hierarchical mass pattern.

The gauge symmetry of the SM as well as those of many extensional models by themselves fix only the gauge bosons. The fermions and Higgs contents have to be chosen somewhat arbitrarily. In the SM, these choices are made in such a way that the neutrinos are massless as mentioned. However, there are other choices based on the SM symmetry that neutrinos become massive. We know these from the popular seesaw [4] and radiative [5] models. Particularly, the models based on the $SU(3)_C \otimes SU(3)_L \otimes U(1)_X$ gauge unification group [6, 7, 8], called 3-3-1 models, give more stricter fermion contents. Indeed, only three fermion generations are acquired as a result of the anomaly cancellation and the condition of QCD asymptotic freedom. The arbitrariness in this case are only behind which SM singlets put in the bottoms of the lepton triplets? In some scenarios, exotic leptons may exist in the singlets. Result of this is quite similar the case of the SM neutrinos. As a fact, the mechanisms of the Zee's type [5] for neutrino masses arise which been explored in Ref.[9].

Forbidding the exotic leptons, there are two main versions of the 3-3-1 models as far as minimal lepton sectors is concerned. In one of them [6] the three known left-handed lepton components for each generation are associated to three $SU(3)_L$ triplets as $(\nu_l, l, l^c)_L$, in which l^c_L is related to the right-handed isospin singlet of the charged lepton l in the SM. No extra leptons are needed and therefore it calls that a minimal 3-3-1 model. In the variant model [7] three $SU(3)_L$ lepton triplets are of the form $(\nu_l, l, \nu_l^c)_L$, where ν_l^c is related to the right-handed component of the neutrino field ν_l , thus called a model with the right-handed neutrinos. This kind of the 3-3-1 models requires only a more economical Higgs sector for breaking the gauge symmetry and generating the fermion masses. Among the new gauge bosons in this model, the neutral non-Hermitian bilepton field X^0 may give promising signature in accelerator experiments and may be also the source of neutrino oscillations [10]. In the current paper, the neutrinos of the 3-3-1 model with right-handed neutrinos is a subject for extended study.

The 3-3-1 model with right-handed neutrinos gives the tree level neutrino mass spectrum with three Dirac fermions, one massless and two degenerate in mass [11]. This is clearly not realistic under the experimental data. However, this pattern may be severely changed by quantum effects and gives rise to an inverted hierarchy mass pattern. This is a *specific* feature of the 3-3-1 model with right-handed neutrinos which was considered in Ref.[11] (see also [12]), but such effects exist in the very high level of the loop corrections.

Some years ago, one of us was proposed the construction of the supersymmetric 3-3-1 model with right-handed neutrino [13]. In this paper, we explore a consistent neutrino mass

spectrum of such a type but in a different side. Namely, that tree level mass spectrum will become a real one of the massive neutrinos due to mixing among them with the neutralinos in the supersymmetric version of the model [13]. In this case, we show that an inverted hierarchy mass spectrum for the neutrinos may be obtained but at only required the tree level contribution which can fit the current data with some minor fine-tuning. Thus, our result differs from many extensions of the SM. As far as the mechanism concerned, it obviously keeps in the kind of an seesaw one. It is not as in the case of the minimal 3-3-1 model, in which its supersymmetric version [14] gives only the real lepton mass spectra when the one loop corrections added [15]. Moreover, in our case the charged leptons always gain consistent masses from different impacts due to mixing among the neutrinos with the neutralinos.

The outline of this work is as follows. In Sec. 2 we review the concept of R-symmetry and R-parity, in order to apply this concept on the supersymmetric 3-3-1 model with right-handed neutrinos. In Sec. 3 we define the R-charge in our model in order to get similar results as in the Minimal Supersymmetric Standard Model (MSSM). While in Sec. 3.1 we impose another discrete symmetry that allow neutrino masses but forbid the proton decay. On Sec. 4 we calculate the fermion masses in our model. Our conclusions are found in the last section. At least, in Appendix A, we present the mass matrix elements of the neutral fermions.

2 R Symmetry

It is important to note that the SM can explain the conservation of lepton number (L) and of baryon number (B) without needing to any discrete symmetry. However, this is not the case of supersymmetric theories where only if interactions of conserving both L and B are required, one has to impose one discrete symmetry. This section recalls how R-parity emerged as a discrete remnant of continuous U(1) R-symmetry which is necessarily broken so that the gauginos and gluinos to acquire masses in the MSSM.

2.1 R Symmetry in Superspace Formalism

The R-symmetry was introduced in 1975 by A. Salam and J. Strathdee [16] and in an independent way by P. Fayet [17] to avoid the interactions that violate either lepton number or baryon number. There is very nice review about this subject in Refs.[18, 19].

The concept of R-symmetry is better understood in superspace formalism, where the R-symmetry is a U(1) continuous symmetry, parametrized by α . The operator which produces this symmetry is going to be denoted as $\mathbf{R}$. This operator acts on the superspace coordinate θ , $\bar{\theta}$ as follows [20]

$$\begin{aligned}\mathbf{R}\theta &\rightarrow e^{-i\alpha}\theta, \\ \mathbf{R}\bar{\theta} &\rightarrow e^{i\alpha}\bar{\theta}.\end{aligned}\tag{2.1}$$

Hence the θ has R-charge to be $R(\theta) = -1$, while $\bar{\theta}$ is $R(\bar{\theta}) = 1$.

The operator $\mathbf{R}$ acts on chiral superfields $\Phi(x, \theta, \bar{\theta})$ and anti-chiral superfields $\bar{\Phi}(x, \theta, \bar{\theta})$, respectively, in the following way [21]

$$\mathbf{R}\Phi(x, \theta, \bar{\theta}) = e^{2in_\Phi\alpha}\Phi(x, e^{-i\alpha}\theta, e^{i\alpha}\bar{\theta}), \quad (2.2)$$

$$\mathbf{R}\bar{\Phi}(x, \theta, \bar{\theta}) = e^{-2in_\Phi\alpha}\bar{\Phi}(x, e^{-i\alpha}\theta, e^{i\alpha}\bar{\theta}), \quad (2.3)$$

where $2n_\Phi$ is the R-charge of the above chiral superfield. This new charge n_Φ is an additive conserved quantum number. This operator acts on the vectorial superfield by the rule

$$\mathbf{R}V(x, \theta, \bar{\theta}) = V(x, e^{-i\alpha}\theta, e^{i\alpha}\bar{\theta}). \quad (2.4)$$

The expansion of the superfields in terms of θ and $\bar{\theta}$, see [21], is given by

$$\begin{aligned} \Phi(x, \theta, \bar{\theta}) = & A(x) + \sqrt{2}\theta\psi(x) + \theta\theta F(x) \\ & + i\theta\sigma^m\bar{\theta}\partial_m A(x) - \frac{i}{\sqrt{2}}(\theta\theta)\partial_m\psi(x)\sigma^m\bar{\theta} \\ & + \frac{1}{4}(\theta\theta)(\bar{\theta}\bar{\theta})\square A(x), \end{aligned} \quad (2.5)$$

$$\begin{aligned} \bar{\Phi}(x, \theta, \bar{\theta}) = & \bar{A}(x) + \sqrt{2}\bar{\theta}\bar{\psi}(x) + \bar{\theta}\bar{\theta}\bar{F}(x) \\ & - i\theta\sigma^m\bar{\theta}\partial_m\bar{A}(x) + \frac{i}{\sqrt{2}}(\bar{\theta}\bar{\theta})\theta\sigma^m\partial_m\bar{\psi}(x) \\ & + \frac{1}{4}(\theta\theta)(\bar{\theta}\bar{\theta})\square\bar{A}(x), \end{aligned} \quad (2.6)$$

$$\begin{aligned} V_{WZ}(x, \theta, \bar{\theta}) = & -\theta\sigma^m\bar{\theta}A_m(x) + i(\theta\theta)\bar{\theta}\bar{\lambda}(x) - i(\bar{\theta}\bar{\theta})\theta\lambda(x) \\ & + \frac{1}{2}(\theta\theta)(\bar{\theta}\bar{\theta})D(x), \text{ expansion in Wess-Zumino gauge,} \end{aligned} \quad (2.7)$$

where $A(x)$, $F(x)$ and $D(x)$ are scalar fields; $\psi(x)$ and $\lambda(x)$ are fermion fields, while $A_m(x)$ is vector field.

Combining Eqs.(2.2) and (2.5) we get the following transformations for the field components, respectively

$$\left. \begin{aligned} A(x) & \xrightarrow{\mathbf{R}} e^{2in_\Phi\alpha}A(x) \\ \psi(x) & \xrightarrow{\mathbf{R}} e^{2i(n_\Phi-\frac{1}{2})\alpha}\psi(x) \\ F(x) & \xrightarrow{\mathbf{R}} e^{2i(n_\Phi-1)\alpha}F(x) \end{aligned} \right\}. \quad (2.8)$$

Similarly, for the anti-chiral superfield we get

$$\left. \begin{aligned} \bar{A}(x) & \xrightarrow{\mathbf{R}} e^{-2in_\Phi\alpha}\bar{A}(x) \\ \bar{\psi}(x) & \xrightarrow{\mathbf{R}} e^{-2i(n_\Phi-\frac{1}{2})\alpha}\bar{\psi}(x) \\ \bar{F}(x) & \xrightarrow{\mathbf{R}} e^{-2i(n_\Phi-1)\alpha}\bar{F}(x) \end{aligned} \right\}. \quad (2.9)$$

From Eqs.(2.4) and (2.7), the field components in the vector superfield transform as

$$\left. \begin{array}{ccc} A_m(x) & \xrightarrow{\mathbf{R}} & A_m(x) \\ \lambda(x) & \xrightarrow{\mathbf{R}} & e^{i\alpha}\lambda(x) \\ \bar{\lambda}(x) & \xrightarrow{\mathbf{R}} & e^{-i\alpha}\bar{\lambda}(x) \\ D(x) & \xrightarrow{\mathbf{R}} & D(x) \end{array} \right\}. \quad (2.10)$$

The transformations in Eqs.(2.8), (2.9) and (2.10) can be rewritten in terms of 4-components spinors as [19]

$$\begin{array}{ccc} A_m(x) & \xrightarrow{\mathbf{R}} & A_m(x), & A(x) & \xrightarrow{\mathbf{R}} & e^{2in_\Phi\alpha} A(x), \\ \Lambda(x) & \xrightarrow{\mathbf{R}} & e^{i\gamma_5\alpha} \Lambda(x), & \bar{A}(x) & \xrightarrow{\mathbf{R}} & e^{-2in_\Phi\alpha} \bar{A}(x), \\ D(x) & \xrightarrow{\mathbf{R}} & D(x), & \Psi(x) & \xrightarrow{\mathbf{R}} & e^{2i\gamma_5(n_\Phi-1/2)\alpha} \Psi(x), \\ & & & F(x) & \xrightarrow{\mathbf{R}} & e^{2i(n_\Phi-1)\alpha} F(x), \\ & & & \bar{F}(x) & \xrightarrow{\mathbf{R}} & e^{-2i(n_\Phi-1)\alpha} \bar{F}(x). \end{array} \quad (2.11)$$

In Eq.(2.11), Λ that is the Majorana spinor represents the gauginos, while $\Psi(x)$ represents the Dirac spinor for quarks and leptons.

For products of left-handed chiral superfields, it is to be noted that

$$\mathbf{R} \prod_a \Phi_a(x, \theta, \bar{\theta}) = e^{2i \sum_a n_a \alpha} \prod_a \Phi_a(x, e^{-i\alpha}\theta, e^{i\alpha}\bar{\theta}).$$

Thus, the general superfield terms given below

$$\begin{aligned} & \int d^4\theta \bar{\Phi}(x, \theta, \bar{\theta}) \Phi(x, \theta, \bar{\theta}), \\ & \int d^4\theta \bar{\Phi}(x, \theta, \bar{\theta}) e^{V(x, \theta, \bar{\theta})} \Phi(x, \theta, \bar{\theta}), \\ & \int d^2\theta \prod_a \Phi_a(x, \theta, \bar{\theta}), \quad \text{if } \sum_a n_a = 1, \end{aligned} \quad (2.12)$$

are all R-invariant.

2.2 Continuous R-Symmetry in MSSM

In the MSSM [22] the left-handed fermions are in doublets, whereas the right-handed antifermions are in singlets: $\hat{L}_L \sim (1, \mathbf{2}, -1)$, $\hat{l}_L^c \sim (1, \mathbf{1}, 2)$ and $\hat{Q}_L \sim (3, \mathbf{2}, 1/3)$, $\hat{u}_L^c \sim (3^*, \mathbf{1}, -4/3)$, $\hat{d}_L^c \sim (3^*, \mathbf{1}, 2/3)$. The Higgs bosons are in doublets, $\hat{H}_1 \sim (1, \mathbf{2}, -1)$ and $\hat{H}_2 \sim (1, \mathbf{2}, 1)$. With these multiplets, the superpotential of the model is written as

$$\begin{aligned} W_2 &= \mu \epsilon \hat{H}_1 \hat{H}_2 + \mu_{0a} \epsilon \hat{L}_{aL} \hat{H}_2, \\ W_3 &= f_{ab}^l \epsilon \hat{L}_{aL} \hat{H}_1 \hat{l}_{bL}^c + f_{ij}^u \epsilon \hat{Q}_{iL} \hat{H}_2 \hat{u}_{jL}^c + f_{ij}^d \epsilon \hat{Q}_{iL} \hat{H}_1 \hat{d}_{jL}^c \\ &\quad + \lambda_{abc} \epsilon \hat{L}_{aL} \hat{L}_{bL} \hat{l}_{cL}^c + \lambda'_{iaj} \epsilon \hat{Q}_{iL} \hat{L}_{aL} \hat{d}_{jL}^c + \lambda''_{ijk} \hat{d}_{iL}^c \hat{u}_{jL}^c \hat{d}_{kL}^c. \end{aligned} \quad (2.13)$$

Hereafter, the superscript L will be removed from the superfields, and the $SU(2)$ indices are default. The superscript c indicates the charge conjugation and ϵ is the antisymmetric $SU(2)$ tensor. The sub-indices a, b, c run over the lepton generations e, μ, τ and $i, j, k = 1, 2, 3$ run over the quark ones.

Because of Eqs.(2.2) and (2.3), the terms proportional to λ, λ' and λ'' are forbidden by the R-symmetry. The following example illustrates this statement. Suppose that

$$\begin{aligned} n_{H_1} &= n_{H_2} = 0, \text{ for } H_1, H_2, \\ n_Q &= n_u = n_d = n_L = n_l = \frac{1}{2}, \text{ for } Q, u^c, d^c, L, l^c, \end{aligned} \quad (2.14)$$

which imply

$$\hat{H}_{1,2}(x, \theta, \bar{\theta}) \xrightarrow{\mathbf{R}} \hat{H}_{1,2}(x, e^{-i\alpha}\theta, e^{i\alpha}\bar{\theta}), \quad (2.15)$$

$$\hat{\Phi}(x, \theta, \bar{\theta}) \xrightarrow{\mathbf{R}} e^{i\alpha} \hat{\Phi}(x, e^{-i\alpha}\theta, e^{i\alpha}\bar{\theta}), \quad \hat{\Phi} = Q, u^c, d^c, L, l^c. \quad (2.16)$$

By Eqs.(2.8) and (2.9), their components transform as

$$\begin{aligned} H_{1,2}(x) &\xrightarrow{\mathbf{R}} H_{1,2}(x), \\ \tilde{H}_{1,2}(x) &\xrightarrow{\mathbf{R}} e^{-i\alpha} \tilde{H}_{1,2}(x), \\ \tilde{f}_L(x) &\xrightarrow{\mathbf{R}} e^{i\alpha} \tilde{f}_L(x), \\ \tilde{f}_L^c(x) &\xrightarrow{\mathbf{R}} e^{-i\alpha} \tilde{f}_L^c(x), \\ \Psi(x) &\xrightarrow{\mathbf{R}} \Psi(x). \end{aligned} \quad (2.17)$$

We recall that $H_{1,2}(x)$ are the Higgs bosons, $\tilde{H}_{1,2}(x)$ are the higgsinos, $\tilde{f}$ are the squarks and sleptons, and $\Psi(x)$ are the quarks and leptons. The consequence of the above transformation is summarized as

$$\begin{aligned} \text{ordinary particle} &\xrightarrow{\mathbf{R}} \text{ordinary particle}, \\ \text{supersymmetric partner} &\xrightarrow{\mathbf{R}} e^{\pm i\alpha} \text{supersymmetric partner}. \end{aligned} \quad (2.18)$$

Under the transformation law in Eq.(2.17), the conserving terms are given by

$$W = f^l \hat{L} \hat{H}_1 \hat{l}^c + f^d \hat{Q} \hat{H}_1 \hat{d}^c + f^u \hat{Q} \hat{H}_2 \hat{u}^c. \quad (2.19)$$

Therefore, the couplings λ, λ' and λ'' are forbidden by the charge assignment given in Eq.(2.14). These terms if they were allowed would induce the rapid proton decay. We allow only the terms from which the fermions in the model gain masses [18].

2.3 Problem with Continuous R-Symmetry, Discrete R-Parity

Because of Eq.(2.12), all the Lagrangians are invariant under the continuous R-symmetry and this obviously avoids the proton decay. However, such an unbroken continuous R-symmetry which acts on the gaugino and gluino mass terms would maintain them massless, even after a spontaneous breaking of the supersymmetry. To see this, let us remember that the gaugino's mass term is given by [23]

$$m_\lambda (\lambda\lambda + \bar{\lambda}\bar{\lambda}), \quad (2.20)$$

which, under the R-symmetry (2.10), transforms as

$$m_\lambda (e^{2i\alpha}\lambda\lambda + e^{-2i\alpha}\bar{\lambda}\bar{\lambda}). \quad (2.21)$$

As a result, the mass term (2.20) is not invariant under the R-symmetry. This fact forces us to abandon the continuous R-symmetry, in favour of the discrete R-symmetry, called *R-parity*. Thereby the R-parity automatically allows gluinos and other gauginos masses.

The discrete R-parity, denoted by $\mathbf{R}_d$, which is able to solve the above problem can be obtained by putting $\alpha = \pi$. Taking this value into account on Eqs.(2.1), (2.2), (2.3) and (2.4) we get the following transformations

$$\begin{aligned} \mathbf{R}_d \theta &\xrightarrow{\mathbf{R}_d} -\theta, \\ \mathbf{R}_d \bar{\theta} &\xrightarrow{\mathbf{R}_d} -\bar{\theta}, \\ \mathbf{R}_d \Phi(x, \theta, \bar{\theta}) &\xrightarrow{\mathbf{R}_d} e^{2in_\Phi\pi} \Phi(x, -\theta, -\bar{\theta}), \\ \mathbf{R}_d \bar{\Phi}(x, \theta, \bar{\theta}) &\xrightarrow{\mathbf{R}_d} e^{-2in_\Phi\pi} \bar{\Phi}(x, -\theta, -\bar{\theta}), \\ \mathbf{R}_d V(x, \theta, \bar{\theta}) &\xrightarrow{\mathbf{R}_d} V(x, -\theta, -\bar{\theta}). \end{aligned} \quad (2.22)$$

It is worth emphasizing that, under this (discrete) transformation law, the terms $\theta\theta$ and $\theta\theta\bar{\theta}\bar{\theta}$ are invariants which is very helpful in further analysis.

Now, under the discrete symmetry, the components of the superfields transform as:

$$\left. \begin{aligned} A(x) &\xrightarrow{\mathbf{R}_d} e^{2in_\Phi\pi} A(x) \\ \psi(x) &\xrightarrow{\mathbf{R}_d} e^{2i(n_\Phi - \frac{1}{2})\pi} \psi(x) \\ F(x) &\xrightarrow{\mathbf{R}_d} e^{2i(n_\Phi - 1)\pi} F(x) \end{aligned} \right\}, \quad (2.23)$$

$$\left. \begin{aligned} A_m(x) &\xrightarrow{\mathbf{R}_d} A_m(x) \\ \lambda(x) &\xrightarrow{\mathbf{R}_d} -\lambda(x) \\ \bar{\lambda}(x) &\xrightarrow{\mathbf{R}_d} -\bar{\lambda}(x) \\ D(x) &\xrightarrow{\mathbf{R}_d} D(x) \end{aligned} \right\}. \quad (2.24)$$

From (2.24), we see that (2.20) is, of course, invariant under the discrete symmetry as mentioned. Moreover, the last term in (2.12) can be redefined by

$$\int d^2\theta \prod_a \Phi_a(x, \theta, \bar{\theta}) , \quad \text{if } \sum_a n_a = 0. \quad (2.25)$$

In the next, we will show that there is a close connection between R-parity and baryon, lepton number conservation laws. Its origin is in our desire to get supersymmetric theories in which B and L could be conserved, and simultaneously, to avoid unwanted exchanges of spin-0 particles.

2.4 Discrete R-Parity in MSSM

Applying the conditions coming from (2.25) on Lagrangians in (2.13) we get the following equations

$$n_{H_1} + n_{H_2} = 0, \quad n_L + n_{H_2} = 0, \quad (2.26)$$

$$n_{H_1} + n_L + n_l = 0, \quad n_{H_1} + n_Q + n_d = 0, \quad (2.27)$$

$$n_{H_2} + n_Q + n_u = 0, \quad n_Q + n_L + n_d = 0, \quad (2.28)$$

$$2n_L + n_l = 0, \quad 2n_d + n_u = 0. \quad (2.29)$$

Unfortunately, not all of these relations can be satisfied simultaneously. Only some of these constraints can be satisfied. For example, choosing

$$\begin{aligned} n_{H_1} &= 0, \quad n_{H_2} = 0, \quad n_L = \frac{1}{2}, \quad n_Q = \frac{1}{2}, \\ n_l &= -\frac{1}{2}, \quad n_u = -\frac{1}{2}, \quad n_d = -\frac{1}{2}, \end{aligned} \quad (2.30)$$

the superfields will transform as

$$\hat{V}(x, \theta, \bar{\theta}) \xrightarrow{\mathbf{R}_d} \hat{V}(x, -\theta, -\bar{\theta}), \quad (2.31)$$

$$\hat{H}_{1,2}(x, \theta, \bar{\theta}) \xrightarrow{\mathbf{R}_d} \hat{H}_{1,2}(x, -\theta, -\bar{\theta}), \quad (2.32)$$

$$\hat{\Phi}(x, \theta, \bar{\theta}) \xrightarrow{\mathbf{R}_d} -\hat{\Phi}(x, -\theta, -\bar{\theta}), \quad \Phi = Q, \quad u^c, \quad d^c, \quad L, \quad l^c. \quad (2.33)$$

In terms of the field components, we obtain

$$\begin{aligned} A_m(x) &\xrightarrow{\mathbf{R}_d} A_m(x), \quad H_{1,2}(x) \xrightarrow{\mathbf{R}_d} H_{1,2}(x), \quad \tilde{f}(x) \xrightarrow{\mathbf{R}_d} -\tilde{f}(x), \\ \Lambda(x) &\xrightarrow{\mathbf{R}_d} -\Lambda(x), \quad \tilde{H}_{1,2}(x) \xrightarrow{\mathbf{R}_d} -\tilde{H}_{1,2}(x), \quad \Psi(x) \xrightarrow{\mathbf{R}_d} \Psi(x). \end{aligned} \quad (2.34)$$

Therefore, the first condition in (2.26), both conditions in (2.27), and again the first condition in (2.28) are satisfied and this does not happen with the remaining conditions. The terms in the superpotential (2.13) which satisfy the rule with the parameters in (2.30) are

$$W = \mu \epsilon \hat{H}_1 \hat{H}_2 + f_{ab}^l \epsilon \hat{L}_a \hat{H}_1 \hat{l}_b^c + f_{ij}^u \epsilon \hat{Q}_i \hat{H}_2 \hat{u}_j^c + f_{ij}^d \epsilon \hat{Q}_i \hat{H}_1 \hat{d}_j^c. \quad (2.35)$$

Their others terms are forbidden which are behind that λ and λ' are kinds of the lepton number violating parameters while λ'' is a type of the baryon number violating parameter.

Equation (2.34) suggests to classify the particles into two types of so called R-even and R-odd. Here the R-even particles ($R_d = +1$) include the gluons, photon, $W^\pm$ and Z

gauge bosons, the quarks, the leptons and the Higgs bosons. Whereas, the R-odd particles ($R_d = -1$) are their superpartners, i.e., the gluinos, neutralinos, charginos, squarks and sleptons. Therefore, R-parity is parity of R-charge of the continuous U(1) R-symmetry and defined by

$$\text{R-parity} = \begin{cases} +1 & \text{for ordinary particles,} \\ -1 & \text{for their superpartners.} \end{cases} \quad (2.36)$$

The above intimate connection between R-parity and baryon number, lepton number conservation laws can be made explicitly by re-expressing (2.36) in terms of the spin S and the matter-parity $(-1)^{3B+L}$ as follows [25]:

$$\text{R-parity} = (-1)^{2S}(-1)^{3B+L}. \quad (2.37)$$

Therefore, all scalar fields ($S = 0$) can be assigned R values

1. Usual scalars: $B = L = 0 \implies R = +1$
2. Sleptons: $B = 0, L = 1 \implies R = -1$
3. Squarks: $B = \frac{1}{3}, L = 0 \implies R = -1$

Analogously for fermions ($S = 1/2$)

1. Gauginos: $B = L = 0 \implies R = -1$
2. Leptons: $B = 0, L = 1 \implies R = +1$
3. Quarks: $B = \frac{1}{3}, L = 0 \implies R = +1$

Because of the gauge bosons as well as all vectorial fields have $B = L = 0$, R-parity = +1. It is to be noted that the above assignment is correct only for the MSSM, where the vector gauge bosons do not carry the lepton number ($L = 0$).

To finish this section, let us note that there will be a lot of other choices of the charges in (2.29) to forbid the fast proton decay [26]. However, all such choices are due to the action of R-symmetry which in a general way can be written as [27]

$$\Phi \longrightarrow e^{2in_\Phi \frac{2\pi}{N}} \Phi. \quad (2.38)$$

Here it is similar to a Z_N symmetry. Among those choices, there is a possibility which allows neutrinos to gain masses. Indeed, choosing

$$\begin{aligned} n_{H_1} &= n_{H_2} = n_L = n_l = 0, \\ n_Q &= \frac{1}{2}, \quad n_u = n_d = -\frac{1}{2}, \end{aligned} \quad (2.39)$$

we get the following transformation of the superfields

$$(Q, u^c, d^c) \rightarrow -(Q, u^c, d^c), (L, l^c, H_1, H_2) \rightarrow (L, l^c, H_1, H_2). \quad (2.40)$$

The terms which are allowed by this new R-parity are obtained by

$$\begin{aligned} W = & \mu \epsilon \hat{H}_1 \hat{H}_2 + \mu_{0a} \hat{L}_a \hat{H}_2 + f_{ab}^l \epsilon \hat{L}_a \hat{H}_1 \hat{l}_b^c + f_{ij}^u \epsilon \hat{Q}_i \hat{H}_2 \hat{u}_j^c + f_{ij}^d \epsilon \hat{Q}_i \hat{H}_1 \hat{d}_j^c \\ & + \lambda_{abc} \epsilon \hat{L}_a \hat{L}_b \hat{l}_c^c + \lambda'_{iaj} \epsilon \hat{Q}_i \hat{L}_a \hat{d}_j^c. \end{aligned} \quad (2.41)$$

As shown in Refs. [28, 29, 30], this superpotential gives neutrinos masses.

3 Discrete R-Parity in SUSY331RN

In the supersymmetric 3-3-1 model with right-handed neutrinos (SUSY331RN) [13], the fermionic content is the following: the left-handed fermions are in triplets/antitriplets $L_a = (\nu_a, l_a, \nu_a^c)_L \sim (\mathbf{1}, \mathbf{3}, -1/3)$, $a = e, \mu, \tau$; $Q_{\alpha L} = (d_\alpha, u_\alpha, d'_\alpha) \sim (\mathbf{3}, \mathbf{3}^*, 0)$, $\alpha = 1, 2$, $Q_{3L} = (u_3, d_3, u') \sim (\mathbf{3}, \mathbf{3}, 1/3)$. The right-handed components are in singlets: $l_a^c \sim (\mathbf{1}, \mathbf{1}, 1)$, u_i^c, d_i^c , $i = 1, 2, 3$, which are similar to those in the SM. In addition, the exotic quarks transform as $u'^c \sim (\mathbf{3}^*, \mathbf{1}, -2/3)$, $d'_\alpha \sim (\mathbf{3}^*, \mathbf{1}, 1/3)$. The scalar content is minimally formed by three Higgs triplets: $\eta = (\eta_1^0, \eta^-, \eta_2^0)^T \sim (\mathbf{1}, \mathbf{3}, -1/3)$; $\chi = (\chi_1^0, \chi^-, \chi_2^0)^T \sim (\mathbf{1}, \mathbf{3}, -1/3)$ and $\rho = (\rho_1^+, \rho^0, \rho_2^+)^T \sim (\mathbf{1}, \mathbf{3}, 2/3)$. The complete set of fields on the 331SUSYRN is given in Ref. [13].

In the model under consideration, the superpotential is given by

$$\begin{aligned} W_2 = & \mu_{0a} \hat{L}_a \hat{\eta}' + \mu_{1a} \hat{L}_a \hat{\chi}' + \mu_\eta \hat{\eta} \hat{\eta}' + \mu_\chi \hat{\chi} \hat{\chi}' + \mu_2 \hat{\eta} \hat{\chi}' + \mu_3 \hat{\chi} \hat{\eta}' + \mu_\rho \hat{\rho} \hat{\rho}', \\ W_3 = & \lambda_{1ab} \hat{L}_a \hat{\rho}' \hat{l}_b^c + \lambda_{2a} \epsilon \hat{L}_a \hat{\chi} \hat{\rho} + \lambda_{3a} \epsilon \hat{L}_a \hat{\eta} \hat{\rho} + \lambda_{4ab} \epsilon \hat{L}_a \hat{L}_b \hat{\rho} + \kappa_{1i} \hat{Q}_3 \hat{\eta}' \hat{u}_i^c \\ & + \kappa'_1 \hat{Q}_3 \hat{\eta}' \hat{u}^c + \kappa_{2i} \hat{Q}_3 \hat{\chi}' \hat{u}_i^c + \kappa'_2 \hat{Q}_3 \hat{\chi}' \hat{u}^c + \kappa_{3\alpha i} \hat{Q}_\alpha \hat{\eta} \hat{d}_i^c \\ & + \kappa'_{3\alpha\beta} \hat{Q}_\alpha \hat{\eta} \hat{d}_\beta^c + \kappa_{4\alpha i} \hat{Q}_\alpha \hat{\rho} \hat{u}_i^c + \kappa'_{4\alpha} \hat{Q}_\alpha \hat{\rho} \hat{u}^c + \kappa_{5i} \hat{Q}_3 \hat{\rho}' \hat{d}_i^c + \kappa'_{5\beta} \hat{Q}_3 \hat{\rho}' \hat{d}_\beta^c \\ & + \kappa_{6\alpha i} \hat{Q}_\alpha \hat{\chi} \hat{d}_i^c + \kappa'_{6\alpha\beta} \hat{Q}_\alpha \hat{\chi} \hat{d}_\beta^c + f_1 \epsilon \hat{\rho} \hat{\chi} \hat{\eta} + f'_1 \epsilon \hat{\rho}' \hat{\chi}' \hat{\eta}' + \zeta_{\alpha\beta\gamma} \epsilon \hat{Q}_\alpha \hat{Q}_\beta \hat{Q}_\gamma \\ & + \lambda'_{\alpha ai} \hat{Q}_\alpha \hat{L}_a \hat{d}_i^c + \lambda''_{ijk} \hat{d}_i^c \hat{u}_j^c \hat{d}_k^c + \xi_{1ij\beta} \hat{d}_i^c \hat{u}_j^c \hat{d}_\beta^c + \xi_{2\alpha\alpha\beta} \hat{Q}_\alpha \hat{L}_a \hat{d}_\beta^c \\ & + \xi_{3i\beta} \hat{d}_i^c \hat{u}^c \hat{d}_\beta^c + \xi_{4ij} \hat{d}_i^c \hat{u}^c \hat{d}_j^c + \xi_{5\alpha i\beta} \hat{d}_\alpha^c \hat{u}_i^c \hat{d}_\beta^c + \xi_{6\alpha\beta} \hat{d}_\alpha^c \hat{u}^c \hat{d}_\beta^c. \end{aligned} \quad (3.1)$$

Applying the conditions coming from (2.25) on (3.1), we get the following equations

$$\begin{aligned} n_L + n_{\eta'} &= 0, \quad n_L + n_{\chi'} = 0, \quad n_\eta + n_{\eta'} = 0, \quad n_\chi + n_{\chi'} = 0, \\ n_\eta + n_{\chi'} &= 0, \quad n_\chi + n_{\eta'} = 0, \quad n_\rho + n_{\rho'} = 0, \\ n_L + n_{\rho'} + n_l &= 0, \quad n_L + n_\chi + n_\rho = 0, \quad n_L + n_\eta + n_\rho = 0, \quad 2n_L + n_\rho = 0, \\ n_{Q_3} + n_{\eta'} + n_u &= 0, \quad n_{Q_3} + n_{\eta'} + n_{u'} = 0, \quad n_{Q_3} + n_{\chi'} + n_u = 0, \\ n_{Q_3} + n_{\chi'} + n_{u'} &= 0, \quad n_{Q_3} + n_{\rho'} + n_d = 0, \quad n_{Q_3} + n_{\rho'} + n_{d'} = 0, \\ n_{Q_\alpha} + n_\eta + n_d &= 0, \quad n_{Q_\alpha} + n_\eta + n_{d'} = 0, \quad n_{Q_\alpha} + n_\chi + n_d = 0, \end{aligned}$$

$$\begin{aligned}
n_{Q_\alpha} + n_\chi + n_{d'} &= 0, \quad n_{Q_\alpha} + n_\rho + n_u = 0, \quad n_{Q_\alpha} + n_\rho + n_{u'} = 0, \\
3n_{Q_\alpha} &= 0, \quad n_\rho + n_\chi + n_\eta = 0, \quad n_{\rho'} + n_{\chi'} + n_{\eta'} = 0, \\
n_{Q_\alpha} + n_L + n_d &= 0, \quad n_{Q_\alpha} + n_L + n_{d'} = 0, \\
2n_d + n_u &= 0, \quad n_d + n_{d'} + n_u = 0, \\
n_d + n_{d'} + n_{u'} &= 0, \quad 2n_d + n_{u'} = 0, \quad 2n_{d'} + n_u = 0, \\
2n_{d'} + n_{u'} &= 0.
\end{aligned} \tag{3.2}$$

Choosing the following R-charges

$$\begin{aligned}
n_\eta &= n_{\eta'} = n_\chi = n_{\chi'} = n_\rho = n_{\rho'} = 0, \\
n_L &= n_{Q_\alpha} = n_{Q_3} = 1/2, \\
n_l &= n_u = n_d = n_{u'} = n_{d'} = -1/2,
\end{aligned} \tag{3.3}$$

and looking at Eq.(2.23), it is easy to see that all the fields $\eta, \eta', \chi, \chi', \rho, \rho', L, Q_\alpha, Q_3, l, u, u', d$ and d' have R-charge equal to one, while their superpartners have opposite R-charge similar to that in the MSSM. The terms which satisfy the defined above symmetry (3.3) are

$$\begin{aligned}
W &= \frac{\mu_\eta}{2} \hat{\eta} \hat{\eta}' + \frac{\mu_\chi}{2} \hat{\chi} \hat{\chi}' + \frac{\mu_\rho}{2} \hat{\rho} \hat{\rho}' + \frac{\mu_2}{2} \hat{\eta} \hat{\chi}' + \frac{\mu_3}{2} \hat{\chi} \hat{\eta}' + \frac{1}{3} \left[\lambda_{1ab} \hat{L}_a \hat{\rho}' \hat{l}_b^c \right. \\
&\quad + \kappa_{1i} \hat{Q}_3 \hat{\eta}' \hat{u}_i^c + \kappa'_1 \hat{Q}_3 \hat{\eta}' \hat{u}'^c + \kappa_{2i} \hat{Q}_3 \hat{\chi}' \hat{u}_i^c + \kappa'_2 \hat{Q}_3 \hat{\chi}' \hat{u}'^c + \kappa_{3\alpha i} \hat{Q}_\alpha \hat{\eta} \hat{d}_i^c \\
&\quad + \kappa'_{3\alpha\beta} \hat{Q}_\alpha \hat{\eta} \hat{d}'^c_\beta + \kappa_{4\alpha i} \hat{Q}_\alpha \hat{\rho} \hat{u}_i^c + \kappa'_{4\alpha} \hat{Q}_\alpha \hat{\rho} \hat{u}'^c + \kappa_{5i} \hat{Q}_3 \hat{\rho}' \hat{d}_i^c + \kappa'_{5\beta} \hat{Q}_3 \hat{\rho}' \hat{d}'^c_\beta \\
&\quad + \kappa_{6\alpha i} \hat{Q}_\alpha \hat{\chi} \hat{d}_i^c + \kappa'_{6\alpha\beta} \hat{Q}_\alpha \hat{\chi} \hat{d}'^c_\beta \\
&\quad \left. + f_1 \epsilon \hat{\rho} \hat{\chi} \hat{\eta} + f'_1 \epsilon \hat{\rho}' \hat{\chi}' \hat{\eta}' \right].
\end{aligned} \tag{3.4}$$

Because of the lepton content of the considering model, the lepton number L obviously does not commute with the gauge symmetry. However, a new conserved charge $\mathcal{L}$ can be constructed through L by making the linear combination $L = x\lambda_3 + y\lambda_8 + \mathcal{L}I$ where λ_3 and λ_8 are the diagonal generators of the $SU(3)_L$ group. Applying this operator on a lepton triplet, the coefficients will be defined

$$L = \frac{2}{\sqrt{3}} \lambda_8 + \mathcal{L}I. \tag{3.5}$$

Moreover, it is useful to produce another conserved charge $\mathcal{B}$ which itself is usual baryon number, $B = \mathcal{B}I$. Thus, the R-parity in this model can be re-expressed via the spin S , new charges $\mathcal{L}$ and $\mathcal{B}$ in terms of

$$\text{R-parity} = (-1)^{2S} (-1)^{3(\mathcal{B}+\mathcal{L})}, \tag{3.6}$$

where the charges $\mathcal{B}$ and $\mathcal{L}$ for the multiplets are defined as follows [11]

<i>Triplet</i>	L	Q_3	χ	η	ρ
$\mathcal{B}$ charge	0	$\frac{1}{3}$	0	0	0
$\mathcal{L}$ charge	$\frac{1}{3}$	$-\frac{2}{3}$	$\frac{4}{3}$	$-\frac{2}{3}$	$-\frac{2}{3}$

(3.7)

<i>Anti – Triplet</i>	Q_α	χ'	η'	ρ'
$\mathcal{B}$ charge	$\frac{1}{3}$	0	0	0
$\mathcal{L}$ charge	$\frac{2}{3}$	$-\frac{4}{3}$	$\frac{2}{3}$	$\frac{2}{3}$

(3.8)

<i>Singlet</i>	l^c	u^c	d^c	u'^c	d'^c
$\mathcal{B}$ charge	0	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$
$\mathcal{L}$ charge	-1	0	0	2	-2

(3.9)

From the superpotential given in Eq.(3.4), it is easy to see that the charged leptons gain mass only from the term

$$-\frac{\lambda_{1ab}}{3}L_a\rho' l_b^c + hc. \quad (3.10)$$

The Higgs fields can have VEVs given by

$$\begin{aligned} \langle \rho \rangle &= (0, u, 0)^T, \quad \langle \rho' \rangle = (0, u', 0)^T, \\ \langle \eta \rangle &= (v, 0, 0)^T, \quad \langle \eta' \rangle = (v', 0, 0)^T, \\ \langle \chi \rangle &= (0, 0, w)^T, \quad \langle \chi' \rangle = (0, 0, w')^T. \end{aligned} \quad (3.11)$$

Combining (3.10) and (3.11) we get mass terms

$$-\frac{\lambda_{1ab}}{3}(l_a l_b^c + \bar{l}_a \bar{l}_b^c)u', \quad (3.12)$$

which lead to the following mass matrix

$$X = \begin{pmatrix} \frac{\lambda_{111}}{3}u' & \frac{\lambda_{112}}{3}u' & \frac{\lambda_{113}}{3}u' \\ \frac{\lambda_{121}}{3}u' & \frac{\lambda_{122}}{3}u' & \frac{\lambda_{123}}{3}u' \\ \frac{\lambda_{131}}{3}u' & \frac{\lambda_{132}}{3}u' & \frac{\lambda_{133}}{3}u' \end{pmatrix}.$$

Hence, all the charged leptons get mass. Notice that only VEV of ρ' is enough to give the charged leptons masses.

Remind that due to conservation of the R -parity defined in Eq.(3.3), there are no terms which give neutrinos masses in the superpotential. Thus, in this case the neutrinos remain massless. From another hand, the gaugino mass terms are given by

$$\mathcal{L}_{GMT} = -\frac{m_\lambda}{2} \left[\sum_{a=1}^8 (\lambda_A^a \lambda_A^a + \bar{\lambda}_A^a \bar{\lambda}_A^a) \right] - \frac{m'}{2} (\lambda_B \lambda_B + \bar{\lambda}_B \bar{\lambda}_B). \quad (3.13)$$

Remind that in this model the non-Hermitian gauge bosons are defined as

$$\begin{aligned} \sqrt{2} W_m^\pm &= V_m^1 \mp iV_m^2, \\ \sqrt{2} Y_m^\pm &= V_m^6 \pm iV_m^7, \\ \sqrt{2} X_m^0 &= V_m^4 - iV_m^5. \end{aligned} \quad (3.14)$$

According to these equations, gauginos of the model are defined as

$$\begin{aligned}\sqrt{2} \lambda_W^\pm &= \lambda_A^1 \mp i\lambda_A^2, \\ \sqrt{2} \lambda_Y^\pm &= \lambda_A^6 \pm i\lambda_A^7, \\ \sqrt{2} \lambda_{X^0} &= \lambda_A^4 - i\lambda_A^5.\end{aligned}\tag{3.15}$$

Then, in terms of these fields, Eq.(3.13) can be rewritten as

$$-m_\lambda(\lambda_W^- \lambda_W^+ + \lambda_Y^- \lambda_Y^+ + \lambda_{X^0} \lambda_{X^{0*}}) - \frac{m_\lambda}{2}(\lambda_A^3 \lambda_A^3 + \lambda_A^8 \lambda_A^8) - \frac{m'}{2} \lambda_B \lambda_B + hc \tag{3.16}$$

The mass matrix of charginos and higgsinos arises from the following Lagrangian

$$\begin{aligned}\mathcal{L}_{H\tilde{H}\tilde{V}} &= -\frac{ig}{\sqrt{2}}[\tilde{\eta}\lambda^a\eta\bar{\lambda}_A^a - \bar{\eta}\lambda^a\tilde{\eta}\lambda_A^a + \tilde{\rho}\lambda^a\rho\bar{\lambda}_A^a - \bar{\rho}\lambda^a\tilde{\rho}\lambda_A^a + \tilde{\chi}\lambda^a\chi\bar{\lambda}_A^a - \bar{\chi}\lambda^a\tilde{\chi}\lambda_A^a \\ &\quad - \tilde{\eta}'\lambda^{*a}\eta'\bar{\lambda}_A^a + \bar{\eta}'\lambda^{*a}\tilde{\eta}'\lambda_A^a - \tilde{\rho}'\lambda^{*a}\rho'\bar{\lambda}_A^a + \bar{\rho}'\lambda^{*a}\tilde{\rho}'\lambda_A^a - \tilde{\chi}'\lambda^{*a}\chi'\bar{\lambda}_A^a + \bar{\chi}'\lambda^{*a}\tilde{\chi}'\lambda_A^a] \\ &\quad - \frac{ig'}{\sqrt{2}}\left[-\frac{1}{3}(\tilde{\eta}\eta\bar{\lambda}_B - \bar{\eta}\tilde{\eta}\lambda_B) - \frac{1}{3}(\tilde{\chi}\chi\bar{\lambda}_B - \bar{\chi}\tilde{\chi}\lambda_B) + \frac{1}{3}(\tilde{\eta}'\eta'\bar{\lambda}_B - \bar{\eta}'\tilde{\eta}'\lambda_B) \right. \\ &\quad \left. + \frac{1}{3}(\tilde{\chi}'\chi'\bar{\lambda}_B - \bar{\chi}'\tilde{\chi}'\lambda_B) + \frac{2}{3}(\tilde{\rho}\rho\bar{\lambda}_B - \bar{\rho}\tilde{\rho}\lambda_B) - \frac{2}{3}(\tilde{\rho}'\rho'\bar{\lambda}_B - \bar{\rho}'\tilde{\rho}'\lambda_B)\right].\end{aligned}\tag{3.17}$$

Using Eq.(3.15), we can rewrite the Lagrangian as

$$\begin{aligned}\mathcal{L}_{H\tilde{H}\tilde{V}} &= +ig[v\tilde{\eta}^-\lambda_W^+ + w\tilde{\chi}^-\lambda_Y^+ + u(\tilde{\rho}_1^+\lambda_W^- + \tilde{\rho}_2^+\lambda_Y^-) - v'\tilde{\eta}'^+\lambda_W^- - w'\tilde{\chi}'^+\lambda_Y^- \\ &\quad - u'(\tilde{\rho}_1'^-\lambda_W^+ + \tilde{\rho}_2'^-\lambda_Y^+)] - \frac{igv}{\sqrt{2}}\tilde{\eta}_1^0\lambda_A^3 - igv\tilde{\eta}_2^0\lambda_{X^0} - \frac{igv}{\sqrt{2}}\tilde{\eta}_1^0\lambda_A^8 \\ &\quad - igw\tilde{\chi}_1^0\lambda_{X^{0*}} + igw\sqrt{\frac{2}{3}}\tilde{\chi}_2^0\lambda_A^8 + \frac{igu}{\sqrt{2}}\tilde{\rho}^0\lambda_A^3 - \frac{igu}{\sqrt{6}}\tilde{\rho}^0\lambda_A^8 + \frac{igv'}{\sqrt{2}}\tilde{\eta}_1'^0\lambda_A^3 + igv'\tilde{\eta}_2'^0\lambda_{X^{0*}} \\ &\quad + \frac{igv'}{\sqrt{6}}\tilde{\eta}_1'^0\lambda_A^8 + igw'\tilde{\chi}_1'^0\lambda_{X^0} - igw'\sqrt{\frac{2}{3}}\tilde{\chi}_2'^0\lambda_A^8 + \frac{igu'}{\sqrt{6}}\tilde{\rho}'^0\lambda_A^8 - \frac{igu'}{\sqrt{2}}\tilde{\rho}'^0\lambda_A^3 - ig'u\sqrt{\frac{2}{3}}\tilde{\rho}^0\lambda_B \\ &\quad + \frac{ig'v}{3\sqrt{2}}\tilde{\eta}_1^0\lambda_B + \frac{ig'w}{3\sqrt{2}}\tilde{\chi}_2^0\lambda_B - \frac{ig'v'}{3\sqrt{2}}\tilde{\eta}_1'^0\lambda_B - \frac{ig'w'}{3\sqrt{2}}\tilde{\chi}_2'^0\lambda_B + ig'u'\sqrt{\frac{2}{3}}\tilde{\rho}^0\lambda_B + hc\end{aligned}\tag{3.18}$$

The mass terms of the higgsinos are given by

$$- \frac{\mu_\eta}{2}\tilde{\eta}\tilde{\eta}' - \frac{\mu_\chi}{2}\tilde{\chi}\tilde{\chi}' - \frac{\mu_\rho}{2}\tilde{\rho}\tilde{\rho}' - \frac{f_1}{3}\epsilon(\tilde{\rho}\tilde{\chi}\eta + \rho\tilde{\chi}\tilde{\eta} + \tilde{\rho}\chi\tilde{\eta}) - \frac{f_1'}{3}\epsilon(\tilde{\rho}'\tilde{\chi}'\eta' + \rho'\tilde{\chi}'\tilde{\eta}' + \tilde{\rho}'\chi'\tilde{\eta}') + hc$$

In the terms of field components, the above expression can be rewritten as

$$\begin{aligned}&- \frac{\mu_\eta}{2}(\tilde{\eta}_1^0\tilde{\eta}_1'^0 + \tilde{\eta}^-\tilde{\eta}'^+ + \tilde{\eta}_2^0\tilde{\eta}_2'^0) - \frac{\mu_\rho}{2}(\tilde{\rho}_1^+\tilde{\rho}_1'^- + \tilde{\rho}^0\tilde{\rho}'^0 + \tilde{\rho}_2^+\tilde{\rho}_2'^-) \\ &- \frac{\mu_\chi}{2}(\tilde{\chi}_1^0\tilde{\chi}_1'^0 + \tilde{\chi}^-\tilde{\chi}'^+ + \tilde{\chi}_2^0\tilde{\chi}_2'^0) - \frac{f_1}{3}\left[(\tilde{\rho}^0\tilde{\chi}_2^0 - \tilde{\rho}_2^+\tilde{\chi}^-)\eta_1^0 + (\tilde{\rho}_1^+\tilde{\chi}^- - \tilde{\rho}^0\tilde{\chi}_1^0)\eta_2^0 \right. \\ &\quad + (\tilde{\chi}_2^0\tilde{\eta}_1^0 - \tilde{\chi}_1^0\tilde{\eta}_2^0)\rho^0 + (\tilde{\rho}_2^+\tilde{\eta}^- - \tilde{\rho}^0\tilde{\eta}_2^0)\chi_1^0 + (\tilde{\rho}^0\tilde{\eta}_1^0 - \tilde{\rho}_1^+\tilde{\eta}^-)\chi_2^0\left] - \frac{f_1'}{3}\left[(\tilde{\rho}'^0\tilde{\chi}_2'^0 - \tilde{\rho}_2'^+\tilde{\chi}'^+)\eta_1'^0 \right. \\ &\quad + (\tilde{\rho}_1'^+\tilde{\chi}'^+ - \tilde{\rho}'^0\tilde{\chi}_1'^0)\eta_2'^0 + (\tilde{\chi}_2'^0\tilde{\eta}_1'^0 - \tilde{\chi}_1'^0\tilde{\eta}_2'^0)\rho'^0 + (\tilde{\rho}_2'^+\tilde{\eta}'^- - \tilde{\rho}'^0\tilde{\eta}_2'^0)\chi_1'^0 \\ &\quad \left. + (\tilde{\rho}'^0\tilde{\eta}_1'^0 - \tilde{\rho}_1'^+\tilde{\eta}'^-)\chi_2'^0\right] + hc\end{aligned}\tag{3.19}$$

The neutral gauginos and higgsinos are mixed by matrix N

$$\tilde{\chi}_\alpha^0 = \sum_{\substack{\beta=\text{gauginos}, \\ \text{higgsinos}}} N_{\alpha\beta} \psi_\beta^0, \quad (3.20)$$

where the $\tilde{\chi}_\alpha^0$ forms fifteen physical Majorana particles. While, the charged gauginos mix with higgsinos via two mixing matrices $C_{\alpha\beta}^\pm$

$$\tilde{\chi}_\alpha^\pm = \sum_{\substack{\beta=\text{gauginos}, \\ \text{higgsinos}}} C_{\alpha\beta}^\pm \psi_\beta^\pm \quad (3.21)$$

to form six physical Dirac particles. These matrices will be presented in the next section.

3.1 The Discrete Symmetry for Proton Stability and Neutrino Masses in SUSY331RN

As before, if we choose the R-charges as follows

$$\begin{aligned} n_L &= n_\eta = n_\chi = \frac{1}{2}, \\ n_{\eta'} &= n_{\chi'} = n_u = n_{u'} = -\frac{1}{2}, \\ n_{Q_3} &= n_{\rho'} = 1, \quad n_d = n_{d'} = -2, \\ n_\rho &= -1, \quad n_{Q_\alpha} = \frac{3}{2}, \quad n_l = -\frac{3}{2}, \end{aligned} \quad (3.22)$$

then the terms under this symmetry are obtained by

$$\begin{aligned} W &= \frac{1}{2} \left(\mu_{0a} \hat{L}_a \hat{\eta}' + \mu_{1a} \hat{L}_a \hat{\chi}' + \mu_\eta \hat{\eta} \hat{\eta}' + \mu_\chi \hat{\chi} \hat{\chi}' + \mu_2 \hat{\eta} \hat{\chi}' + \mu_3 \hat{\chi} \hat{\eta}' + \mu_\rho \hat{\rho} \hat{\rho}' \right) \\ &+ \frac{1}{3} \left(\lambda_{1ab} \hat{L}_a \hat{\rho}' \hat{l}_b^c + \lambda_{2a} \epsilon \hat{L}_a \hat{\chi} \hat{\rho} + \lambda_{3a} \epsilon \hat{L}_a \hat{\eta} \hat{\rho} + \lambda_{4ab} \epsilon \hat{L}_a \hat{L}_b \hat{\rho} + \kappa_{1i} \hat{Q}_3 \hat{\eta}' \hat{u}_i^c + \kappa'_1 \hat{Q}_3 \hat{\eta}' \hat{u}'^c \right. \\ &+ \kappa_{2i} \hat{Q}_3 \hat{\chi}' \hat{u}_i^c + \kappa'_2 \hat{Q}_3 \hat{\chi}' \hat{u}'^c + \kappa_{3\alpha i} \hat{Q}_\alpha \hat{\eta} \hat{d}_i^c + \kappa'_{3\alpha\beta} \hat{Q}_\alpha \hat{\eta} \hat{d}_\beta'^c + \kappa_{4\alpha i} \hat{Q}_\alpha \hat{\rho} \hat{u}_i^c + \kappa'_{4\alpha} \hat{Q}_\alpha \hat{\rho} \hat{u}'^c \\ &+ \kappa_{5i} \hat{Q}_3 \hat{\rho}' \hat{d}_i^c + \kappa'_{5\beta} \hat{Q}_3 \hat{\rho}' \hat{d}_\beta^c + \kappa_{6\alpha i} \hat{Q}_\alpha \hat{\chi} \hat{d}_i^c + \kappa'_{6\alpha\beta} \hat{Q}_\alpha \hat{\chi} \hat{d}_\beta'^c + f_1 \epsilon \hat{\rho} \hat{\chi} \hat{\eta} + f'_1 \epsilon \hat{\rho}' \hat{\chi}' \hat{\eta}' \\ &\left. + \lambda'_{\alpha ai} \hat{Q}_\alpha \hat{L}_a \hat{d}_i^c + \xi_{2\alpha a\beta} \hat{Q}_\alpha \hat{L}_a \hat{d}_\beta'^c \right). \end{aligned} \quad (3.23)$$

In this case, it is easy to see that the fields L , l , Q_α , u , u' , $\tilde{\eta}$, $\tilde{\eta}'$, $\tilde{\chi}$, $\tilde{\chi}'$, ρ , ρ' , $\tilde{Q}_3$, $\tilde{d}$ and $\tilde{d}'$ have R -charge equal to one, while the others fields have opposite R -charge.

The superpotential in (3.23) provides us the mass terms for leptons and higgsinos

$$-\frac{\mu_{0a}}{2} L_a \tilde{\eta}' - \frac{\mu_{1a}}{2} L_a \tilde{\chi}' - \frac{\mu_2}{2} \tilde{\eta} \tilde{\chi}' - \frac{\mu_3}{2} \tilde{\chi} \tilde{\eta}' - \frac{\lambda_{2a}}{3} (L_a \tilde{\chi} \rho + \tilde{\rho} L_a \chi) - \frac{\lambda_{3a}}{3} (L_a \tilde{\eta} \rho + \tilde{\rho} L_a \eta) - \frac{\lambda_{4ab}}{3} L_a L_b \rho + hc, \quad (3.24)$$

which in terms of field components get a form

$$\begin{aligned}
& - \frac{\mu_{0a}}{2}(\nu_a \tilde{\eta}_1^{\prime 0} + l_a \tilde{\eta}^{\prime +} + \nu_a^c \tilde{\eta}_2^{\prime 0}) - \frac{\mu_{1a}}{2}(\nu_a \tilde{\chi}_1^{\prime 0} + l_a \tilde{\chi}^{\prime +} + \nu_a^c \tilde{\chi}_2^{\prime 0}) \\
& - \frac{\mu_2}{2}(\tilde{\eta}_1^0 \tilde{\chi}_1^{\prime 0} + \tilde{\eta}^- \tilde{\chi}^{\prime +} + \tilde{\eta}_2^0 \tilde{\chi}_2^{\prime 0}) - \frac{\mu_3}{2}(\tilde{\chi}_1^0 \tilde{\eta}_1^{\prime 0} + \tilde{\chi}^- \tilde{\eta}^{\prime +} + \tilde{\chi}_2^0 \tilde{\eta}_2^{\prime 0}) - \frac{\lambda_{2a}}{3}[(\nu_a^c \tilde{\chi}_1^0 - \nu_a \tilde{\chi}_2^0)\rho^0 \\
& + (\tilde{\rho}^0 \nu_a^c - \tilde{\rho}_2^+ l_a)\chi_1^0 + (\tilde{\rho}_1^+ l_a - \tilde{\rho}^0 \nu_a)\chi_2^0] - \frac{\lambda_{3a}}{3}[(\nu_a^c \tilde{\eta}_1^0 - \nu_a \tilde{\eta}_2^0)\rho^0 + (\tilde{\rho}^0 \nu_a^c - \tilde{\rho}_2^+ l_a)\eta_1^0 \\
& + (\tilde{\rho}_1^+ l_a - \tilde{\rho}^0 \nu_a)\eta_2^0] - \frac{\lambda_{4ab}}{3}(\nu_a^c \nu_b - \nu_b^c \nu_a)\rho^0 + h.c
\end{aligned} \tag{3.25}$$

It is easy to see that the above Lagrangian gives mass terms for neutrinos and forbids the proton decay.

4 Fermion masses

With the above mass terms, we get the mass matrices for the neutral and charged fermions, respectively. Diagonalizing these matrices, we get the physical masses for the fermions.

4.1 Masses of the neutral fermions

Mass Lagrangian for the neutral fermions are easily obtained by

$$\begin{aligned}
\mathcal{M}_{neutral} = & -\frac{1}{2} \left[\mu_{0a}(\nu_{aL} \tilde{\eta}_1^{\prime 0} + \nu_{aL}^c \tilde{\eta}_2^{\prime 0}) + \mu_{1a}(\nu_{aL} \tilde{\chi}_1^{\prime 0} + \nu_{aL}^c \tilde{\chi}_2^{\prime 0}) + \mu_\eta(\tilde{\eta}_1^0 \tilde{\eta}_1^{\prime 0} + \tilde{\eta}_2^0 \tilde{\eta}_2^{\prime 0}) \right. \\
& + \mu_\chi(\tilde{\chi}_1^0 \tilde{\chi}_1^{\prime 0} + \tilde{\chi}_2^0 \tilde{\chi}_2^{\prime 0}) + \mu_\rho \tilde{\rho}^0 \tilde{\rho}^{\prime 0} + \mu_2(\tilde{\eta}_1^0 \tilde{\chi}_1^{\prime 0} + \tilde{\eta}_2^0 \tilde{\chi}_2^{\prime 0}) + \mu_3(\tilde{\chi}_1^0 \tilde{\eta}_1^{\prime 0} + \tilde{\chi}_2^0 \tilde{\eta}_2^{\prime 0}) \Big] \\
& - \frac{1}{3} \left[\lambda_{2a} u(\nu_{aL}^c \tilde{\chi}_1^0 - \nu_{aL} \tilde{\chi}_2^0) - \lambda_{2a} w \nu_{aL} \tilde{\rho}^0 + \lambda_{3a} u(\nu_{aL}^c \tilde{\eta}_1^0 - \nu_{aL} \tilde{\eta}_2^0) \right. \\
& + \lambda_{3a} v \nu_{aL}^c \tilde{\rho}^0 + \lambda_{4ab} u(\nu_{aL}^c \nu_{bL} - \nu_{bL}^c \nu_{aL}) + f_1 v \tilde{\rho}^0 \tilde{\chi}_2^0 + f_1 w \tilde{\rho}^0 \tilde{\eta}_1^0 \\
& + f_1 u(\tilde{\chi}_2^0 \tilde{\eta}_1^0 - \tilde{\chi}_1^0 \tilde{\eta}_2^0) + f_1' v' \tilde{\rho}^{\prime 0} \tilde{\chi}_2^0 + f_1' w' \tilde{\rho}^{\prime 0} \tilde{\eta}_1^0 + f_1' u'(\tilde{\chi}_2^{\prime 0} \tilde{\eta}_1^0 - \tilde{\chi}_1^{\prime 0} \tilde{\eta}_2^0) \Big] \\
& - \frac{m_\lambda}{2}(2\lambda_{X^0} \lambda_{X^{0*}} + \lambda_A^3 \lambda_A^3 + \lambda_A^8 \lambda_A^8) - \frac{m'}{2} \lambda_B \lambda_B \\
& + \frac{igv}{\sqrt{2}} \tilde{\eta}_1^0 \lambda_A^3 + igv \tilde{\eta}_2^0 \lambda_{X^0} + \frac{igv}{\sqrt{6}} \tilde{\eta}_1^0 \lambda_A^8 + igw \tilde{\chi}_1^0 \lambda_{X^{0*}} \\
& - igw \sqrt{\frac{2}{3}} \tilde{\chi}_2^0 \lambda_A^8 - \frac{igu}{\sqrt{2}} \tilde{\rho}^0 \lambda_A^3 + \frac{igu}{\sqrt{6}} \tilde{\rho}^0 \lambda_A^8 - \frac{igv'}{\sqrt{2}} \tilde{\eta}_1^{\prime 0} \lambda_A^3 - igv' \tilde{\eta}_2^{\prime 0} \lambda_{X^{0*}} \\
& - \frac{igv'}{\sqrt{6}} \tilde{\eta}_1^{\prime 0} \lambda_A^8 - igw' \tilde{\chi}_1^{\prime 0} \lambda_{X^0} + igw' \sqrt{\frac{2}{3}} \tilde{\chi}_2^{\prime 0} \lambda_A^8 - \frac{igu'}{\sqrt{6}} \tilde{\rho}^{\prime 0} \lambda_A^8 - \frac{igu'}{\sqrt{2}} \tilde{\rho}^{\prime 0} \lambda_A^3 \\
& + ig'u \sqrt{\frac{2}{3}} \tilde{\rho}^0 \lambda_B - \frac{ig'v}{3\sqrt{2}} \tilde{\eta}_1^0 \lambda_B - \frac{ig'w}{3\sqrt{2}} \tilde{\chi}_2^0 \lambda_B + \frac{ig'v'}{3\sqrt{2}} \tilde{\eta}_1^{\prime 0} \lambda_B + \frac{ig'w'}{3\sqrt{2}} \tilde{\chi}_2^{\prime 0} \lambda_B
\end{aligned}$$

$$-ig'u'\sqrt{\frac{2}{3}}\tilde{\rho}^0\lambda_B + h.c.$$

In the basis Ψ^0 of the form

$$\left(\nu_1 \ \nu_2 \ \nu_3 \ \nu_1^c \ \nu_2^c \ \nu_3^c - i\lambda_A^3 - i\lambda_{X^0} - i\lambda_{X^{0*}} - i\lambda_A^8 - i\lambda_B \ \tilde{\eta}_1^0 \ \tilde{\eta}_1'^0 \ \tilde{\eta}_2^0 \ \tilde{\eta}_2'^0 \ \tilde{\chi}_1^0 \ \tilde{\chi}_1'^0 \ \tilde{\chi}_2^0 \ \tilde{\chi}_2'^0 \ \tilde{\rho}^0 \ \tilde{\rho}'^0\right),$$

the mass Lagrangian can be written as follows

$$-\frac{1}{2}(\Psi^0)^T Y^0 \Psi^0 + h.c. \quad (4.1)$$

Here Y^0 is symmetric matrix with the nonzero elements given in Appendix.A, where the mass eigenstates are given by

$$\tilde{\chi}_i^0 = N_{ij}\Psi_j^0, \quad j = 1, \dots, 21. \quad (4.2)$$

The mass matrix of the neutral fermions consists of three parts: (a) The first part M_ν is the 6×6 mass matrix of the neutrinos which belongs to the 3-3-1 model with right-handed neutrinos; (b) While, the second part M_N is the 15×15 mass matrix of the neutralinos which exists only in its supersymmetric version; (c) The last part $M_{\nu N}$ arises due to mixing among the neutrinos and neutralinos. Thus, the mass matrix for the neutral fermions is signified as follows

$$Y^0 = \begin{pmatrix} M_\nu & M_{\nu N} \\ M_{\nu N}^T & M_N \end{pmatrix}. \quad (4.3)$$

To keep consistent with the low-energy effective theories, some Yukawa couplings in the sub-matrices will be fixed in terms of fine-tunings needed in this model. First, we know that the mass matrix M_ν gives three Dirac eigenstates. Two of them have degenerate eigenvalues $m_\nu = \frac{2u}{3}\sqrt{\lambda_{412}^2 + \lambda_{413}^2 + \lambda_{423}^2}$ and the other one massless. Thus, it is easy to identify the mass splitting m_ν as the value of measured atmospheric neutrino mass difference $\Delta m_{atm} \sim 5 \times 10^{-2}$ eV. By putting $\lambda_{412} = \lambda_{413} = \lambda_{423}$, we get

$$\begin{aligned} \lambda_{412} &= \lambda_{413} = \lambda_{423} = 2 \times 10^{-13}, \\ \lambda_{421} &= \lambda_{431} = \lambda_{432} = -2 \times 10^{-13}. \end{aligned} \quad (4.4)$$

It is worth emphasizing that when the mixing terms turn on, this inverted spectrum will not only give rise to mass splitting between the two degenerate Dirac states, it will also split each Dirac pairs into two non-degenerate Majorana states, resulting in the spectrum with six Majorana eigenstates with four heavier ones and two light ones. Here we are assuming that the solar oscillation is between the two heavier Majorana states.

Finally, to keep the mass constraints from astrophysics and cosmology [31] as well as to be consistent with all the earlier analyses [15], the parameters in the mass matrix M_N can be chosen as a typical example by:

$$\begin{aligned} \mu_\eta &= 300 \text{ GeV}, \quad \mu_\rho = 500 \text{ GeV}, \quad \mu_\chi = 700 \text{ GeV}, \\ \mu_2 &= 50 \text{ GeV}, \quad \mu_3 = 200 \text{ GeV}, \quad m' = 2000 \text{ GeV}, \\ \mu_\lambda &= 3000 \text{ GeV}, \quad f_1 = 1.8, \quad f'_1 = 10^{-3}. \end{aligned} \quad (4.5)$$

Here in this model, the Higgs bosons' VEVs are fixed as follows

$$\begin{aligned} v_\eta &= 15 \text{ GeV}, \quad v_{\eta'} = 10 \text{ GeV}, \\ v_\rho &= 244.9 \text{ GeV}, \quad v_{\rho'} = 13 \text{ GeV}, \\ v_{\chi_2} &= v_{\chi'_2} = 1000 \text{ GeV}, \end{aligned} \quad (4.6)$$

and the value of g given in [31].

Now, the mixing terms rise to correct the effective ones, thus, in some ways we can get the physical masses. In the first case, the parameters in the mixing matrix $M_{\nu N}$ can be chosen as follows

1. For the dimensionless parameters:

$$\begin{aligned} \lambda_{21} &= 0, \quad \lambda_{22} = 0, \quad \lambda_{23} = 0, \\ \lambda_{31} &= 0, \quad \lambda_{32} = 0, \quad \lambda_{33} = 0, \end{aligned} \quad (4.7)$$

2. For the mass-scale parameters (in GeV):

$$\begin{aligned} \mu_{01} &= 10^{-4}, \quad \mu_{02} = 10^{-4}, \quad \mu_{03} = 0, \\ \mu_{11} &= 10^{-5}, \quad \mu_{12} = 0, \quad \mu_{13} = 0. \end{aligned} \quad (4.8)$$

Hence, the neutralinos obtain the masses in GeV as follows

$$\begin{aligned} &3210.2199, \quad 3152.6352, \quad 3152.6352, \quad 3004.7008, \quad 2087.1869, \\ &693.1140, \quad 672.9818, \quad 367.2895, \quad 281.0245, \quad 269.3481, \\ &151.7631, \quad 115.9158, \quad 48.4871, \quad 46.7180, \quad 39.5593. \end{aligned} \quad (4.9)$$

The tauon neutrino gains masses in eV by

$$5.39600 \times 10^{-5}, \quad 4.12596 \times 10^{-7}, \quad (4.10)$$

while the masses of the electron and muon neutrinos are (in eV)

$$\begin{aligned} &6.31927 \times 10^{-2}, \quad 5.49914 \times 10^{-2}, \\ &5.46530 \times 10^{-2}, \quad 4.70848 \times 10^{-2}. \end{aligned} \quad (4.11)$$

In the second case, the values of the parameters in the mixing matrix $M_{\nu N}$ are given in another way as below

1. For the mass-scale parameters:

$$\begin{aligned} \mu_{01} &= 0, \quad \mu_{02} = 0, \quad \mu_{03} = 0, \\ \mu_{11} &= 0, \quad \mu_{12} = 0, \quad \mu_{13} = 0, \end{aligned} \quad (4.12)$$




Figure 1: The inverted hierarchy mass pattern of neutrinos in the model.

2. For the dimensionless parameters:

$$\begin{aligned}\lambda_{21} &= 10^{-6}, \lambda_{22} = 10^{-6}, \lambda_{23} = 10^{-6}, \\ \lambda_{31} &= 10^{-6}, \lambda_{32} = 0, \lambda_{33} = 0.\end{aligned}\tag{4.13}$$

The masses (in GeV) of the neutralinos are

$$\begin{aligned}3210.2199, 3152.6352, 3152.6352, 3004.7008, 2087.1869, \\ 693.1140, 672.9818, 367.2895, 281.0245, 269.3481, \\ 151.7631, 115.9158, 48.4871, 46.7180, 39.5593.\end{aligned}\tag{4.14}$$

The tauon neutrino in this case gains masses in eV by

$$1.95303 \times 10^{-4}, 4.87078 \times 10^{-5}.\tag{4.15}$$

Also, the electron and muon neutrinos have masses in eV

$$\begin{aligned}8.37329 \times 10^{-2}, 5.52047 \times 10^{-2}, \\ 5.51605 \times 10^{-2}, 3.69629 \times 10^{-2}.\end{aligned}\tag{4.16}$$

Notice that the coupling constant g' , λ_{4ab} and the parameter m' appear only in the mass matrix of the neutral fermions, while, those of the charged sector are λ_{1ab} .

As above, we give two typical examples of the values of the mixing terms which not only give the consistent mass spectra of the neutrinos but also keep an large enough hierarchy so that the neutralinos gain the masses satisfying the lower mass limit (> 32.5 GeV) from astrophysics and cosmology. Consequently, the neutrinos in this model yield inverted hierarchy mass patterns as shown in Fig.(1).

4.2 Masses of the charged fermions

The terms contributing to the masses of the charged fermions are

$$\begin{aligned}
\mathcal{M}_{charged} = & -\frac{\lambda_{1ab}}{3}l_a l_b^c u' - \frac{f_1}{3}(\tilde{\rho}_2^+ \tilde{\chi}^- v + \tilde{\rho}_1^+ \tilde{\eta}^- w) - \frac{f'_1}{3}(\tilde{\rho}_2'^- \tilde{\chi}'^+ v' + \tilde{\rho}_1'^- \tilde{\eta}'^+ w') \\
& -m_\lambda(\lambda_W^- \lambda_W^+ + \lambda_Y^- \lambda_Y^+) + ig[v\tilde{\eta}^- \lambda_W^+ + w\tilde{\chi}^- \lambda_Y^+ + u(\tilde{\rho}_1^+ \lambda_W^- + \tilde{\rho}_2^+ \lambda_Y^-) - v'\tilde{\eta}'^+ \lambda_W^- \\
& -w'\tilde{\chi}'^+ \lambda_Y^- - u'(\tilde{\rho}_1'^- \lambda_W^+ + \tilde{\rho}_2'^- \lambda_Y^+)] - \frac{\mu_\eta}{2}\tilde{\eta}^- \tilde{\eta}'^+ - \frac{\mu_\rho}{2}(\tilde{\rho}_1^+ \tilde{\rho}_1'^- + \tilde{\rho}_2^+ \tilde{\rho}_2'^-) - \frac{\mu_\chi}{2}\tilde{\chi}^- \tilde{\chi}'^+ \\
& -\frac{\mu_{0a}}{2}l_a \tilde{\eta}'^+ - \frac{\mu_{1a}}{2}l_a \tilde{\chi}'^+ - \frac{\mu_2}{2}\tilde{\eta}^- \tilde{\chi}'^+ - \frac{\mu_3}{2}\tilde{\chi}^- \tilde{\eta}'^+ - \frac{\lambda_{2a}}{3}l_a \tilde{\rho}_1^+ w + \frac{\lambda_{3a}}{3}l_a \tilde{\rho}_2^+ v + h.c.
\end{aligned}$$

To write the mass matrix, we will choose the following bases

$$\begin{aligned}
\psi^- &= \begin{pmatrix} l_1 & l_2 & l_3 & -i\lambda_W^- & -i\lambda_Y^- & \tilde{\eta}^- & \tilde{\chi}^- & \tilde{\rho}_1'^- & \tilde{\rho}_2'^- \end{pmatrix}^T \\
\psi^+ &= \begin{pmatrix} l_1^c & l_2^c & l_3^c & i\lambda_W^+ & i\lambda_Y^+ & \tilde{\eta}'^+ & \tilde{\chi}'^+ & \tilde{\rho}_1^+ & \tilde{\rho}_2^+ \end{pmatrix}^T
\end{aligned} \tag{4.17}$$

and define

$$\Psi^\pm = \begin{pmatrix} \psi^+ & \psi^- \end{pmatrix}^T. \tag{4.18}$$

With these definitions, the mass term is written in the form

$$-\frac{1}{2}(\Psi^\pm)^T Y^\pm \Psi^\pm + h.c., \tag{4.19}$$

where

$$Y^\pm = \begin{pmatrix} 0 & X^T \\ X & 0 \end{pmatrix}. \tag{4.20}$$

Then, the X matrix is given by

$$X = \begin{pmatrix} \frac{\lambda_{111}}{3}u' & \frac{\lambda_{112}}{3}u' & \frac{\lambda_{113}}{3}u' & 0 & 0 & \frac{1}{2}\mu_{01} & \frac{1}{2}\mu_{11} & \frac{\lambda_{21}}{3}w & -\frac{\lambda_{31}}{3}v \\ \frac{\lambda_{121}}{3}u' & \frac{\lambda_{122}}{3}u' & \frac{\lambda_{123}}{3}u' & 0 & 0 & \frac{1}{2}\mu_{02} & \frac{1}{2}\mu_{12} & \frac{\lambda_{22}}{3}w & -\frac{\lambda_{32}}{3}v \\ \frac{\lambda_{131}}{3}u' & \frac{\lambda_{132}}{3}u' & \frac{\lambda_{133}}{3}u' & 0 & 0 & \frac{1}{2}\mu_{03} & \frac{1}{2}\mu_{13} & \frac{\lambda_{23}}{3}w & -\frac{\lambda_{33}}{3}v \\ 0 & 0 & 0 & -m_\lambda & 0 & -gv' & 0 & gu & 0 \\ 0 & 0 & 0 & 0 & -m_\lambda & 0 & -gw' & 0 & gu \\ 0 & 0 & 0 & gv & 0 & \frac{\mu_\eta}{2} & \frac{\mu_2}{2} & \frac{f_1}{3}w & 0 \\ 0 & 0 & 0 & 0 & gw & \frac{\mu_3}{2} & \frac{\mu_\chi}{2} & 0 & \frac{f_1}{3}v \\ 0 & 0 & 0 & -gu' & 0 & \frac{f'_1}{3}w' & 0 & \frac{\mu_\rho}{2} & 0 \\ 0 & 0 & 0 & 0 & -gu' & 0 & \frac{f'_1}{3}v' & 0 & \frac{\mu_\rho}{2} \end{pmatrix}.$$

The chargino mass matrix $Y^\pm$ is diagonalized by using two unitary matrices, D and E , defined by

$$\tilde{\chi}_i^+ = D_{ij}\Psi_j^+, \quad \tilde{\chi}_i^- = E_{ij}\Psi_j^-, \quad i, j = 1, \dots, 9. \tag{4.21}$$

The characteristic equation for the matrix $Y^\pm$ is

$$\det(Y^\pm - \lambda I) = \det \left[\begin{pmatrix} -\lambda & X^T \\ X & -\lambda \end{pmatrix} \right] = \det(\lambda^2 - X^T \cdot X). \quad (4.22)$$

Since $X^T \cdot X$ is a symmetric matrix, λ^2 must be real and positive because $Y^\pm$ is also symmetric. Hence, to obtain eigenvalues, one only have to calculate $X^T \cdot X$. Then we can write the diagonal mass matrix as

$$M_{SCM} = E^* X D^{-1}. \quad (4.23)$$

To determine E and D , it is useful the following observation

$$M_{SCM}^2 = D X^T \cdot X D^{-1} = E^* X \cdot X^T (E^*)^{-1}, \quad (4.24)$$

which means that D diagonalizes $X^T \cdot X$, while E^* diagonalizes $X \cdot X^T$. In this case we can define the following Dirac spinors:

$$\Psi(\tilde{\chi}_i^+) = \begin{pmatrix} \tilde{\chi}_i^+ & \bar{\tilde{\chi}}_i^- \end{pmatrix}^T, \quad \Psi^c(\tilde{\chi}_i^-) = \begin{pmatrix} \tilde{\chi}_i^- & \bar{\tilde{\chi}}_i^+ \end{pmatrix}^T. \quad (4.25)$$

where $\tilde{\chi}_i^+$ is the particle and $\tilde{\chi}_i^-$ is the anti-particle [22, 32].

Now to get mass values, all the parameters in the neutral sector should be kept in this sector of the charged fermions. Corresponding to the first case in the neutral sector, we have obtained the following masses (in GeV) for the charginos:

$$3156.1474, 3004.4371, 665.1171, 263.3014, 209.0726, 45.9104. \quad (4.26)$$

The ordinary leptons gain masses in GeV as $m_\tau = 1.7766$, $m_\mu = 0.1057$, $m_e = 0.00051$. In this case, the masses have been obtained by using the remaining set of the dimensionless parameters:

$$\begin{aligned} \lambda_{111} &= 1.18 \times 10^{-4}, & \lambda_{112} &= 10^{-7}, & \lambda_{113} &= 10^{-7}, \\ \lambda_{121} &= 10^{-7}, & \lambda_{122} &= 2.44 \times 10^{-2}, & \lambda_{123} &= 10^{-7}, \\ \lambda_{131} &= 10^{-7}, & \lambda_{132} &= 10^{-7}, & \lambda_{133} &= 4.10 \times 10^{-1}. \end{aligned} \quad (4.27)$$

The second case is obtained by changing only the mixing terms as in the neutral sector. As a result, the masses are *the same* as in the previous case. Thus, the ordinary charged leptons get the consistent masses; and, the lightest chargino with the mass of 45.9104 GeV is in the experimental lower limit of 45 GeV. It was shown in this section that, there are nine fermions. However, as mentioned above, by the conservation of R-parity, there are only six charginos.

To summarize, as above we have given at the tree level the consistent masses for the charged leptons and the neutrinos in the supersymmetric 3-3-1 model with right-handed neutrinos. Such a model for the leptons is more simpler than that of the supersymmetric minimal 3-3-1 model [14], where the loop corrections are needed for the masses of the leptons [15]. The charginos and neutralinos in this model gain the masses respectively very smaller than those of the supersymmetric minimal 3-3-1 model [15]. Contrasting with the supersymmetric minimal 3-3-1 model, the MSSM neutralinos and charginos in this model can be directly identified via the mass spectra given above, where the mass constraints on the MSSM particles can be found in Ref.[34].

5 Conclusion

In this article, we have found, in framework of the supersymmetric 3-3-1 model with right-handed neutrinos, definitions of the R charges which are similar with that in the MSSM. This means that in the considered model, there is one discrete symmetry which allows the neutral and charged fermions, gauge bosons and scalar fields to get masses and at same time forbidding the proton decay. Thus, in this case there exists one phenomenology similar to the MSSM with the famous missing transverse energy events, which is specific of the R -parity conservation [33].

We have showed that there is one symmetry which gives neutrinos masses but forbids the proton decay. Unlike the cases with the MSSM and the minimal 3-3-1 model [15], in this model, all the fermions get masses at the tree level.

The famous relation for the R -parity in the MSSM has been generalized to this kind of the 3-3-1 models. In this case it relates to the new conserved charge $\mathcal{L}$. A simple mechanism for the mass generation of the neutrinos has been explored. We have showed that the model naturally gives rise to the neutrinos an inverted hierarchy mass pattern. Moreover, the MSSM superpartners in this model can be explicitly identified which is unlike in the case of supersymmetric extension of the minimal 3-3-1 model.

Acknowledgments

M. C. R would like to thank Vietnam Academy of Science and Technology by the nice hospitality, warm atmosphere during his stay at Institute of Physics to start this work.

A Mass matrix elements of Y^0

The nonzero elements of Y^0 are

$$\begin{aligned}
m_{1,5} &= m_{5,1} = -\frac{u}{3}(\lambda_{421} - \lambda_{412}), \quad m_{1,6} = m_{6,1} = -\frac{u}{3}(\lambda_{431} - \lambda_{413}), \\
m_{1,13} &= m_{13,1} = \frac{\mu_{01}}{2}, \quad m_{1,14} = m_{14,1} = -\frac{\lambda_{31}}{3}u, \quad m_{1,17} = m_{17,1} = \frac{\mu_{11}}{2}, \\
m_{1,18} &= m_{18,1} = -\frac{\lambda_{21}}{3}u, \quad m_{1,20} = m_{20,1} = -\frac{\lambda_{21}}{3}w, \\
m_{2,4} &= m_{4,2} = -\frac{u}{3}(\lambda_{412} - \lambda_{421}), \quad m_{2,6} = m_{6,2} = -\frac{u}{3}(\lambda_{432} - \lambda_{423}), \\
m_{2,13} &= m_{13,2} = \frac{\mu_{02}}{2}, \quad m_{2,14} = m_{14,2} = -\frac{\lambda_{32}}{3}u, \quad m_{2,17} = m_{17,2} = \frac{\mu_{12}}{2}, \\
m_{2,18} &= m_{18,2} = -\frac{\lambda_{22}}{3}u, \quad m_{2,20} = m_{20,2} = -\frac{\lambda_{22}}{3}w, \\
m_{3,4} &= m_{4,3} = -\frac{u}{3}(\lambda_{413} - \lambda_{431}), \quad m_{3,5} = m_{5,3} = -\frac{u}{3}(\lambda_{423} - \lambda_{432}),
\end{aligned}$$

$$\begin{aligned}
m_{3,13} &= m_{13,3} = \frac{\mu_{03}}{2}, \quad m_{3,14} = m_{14,3} = -\frac{\lambda_{33}}{3}u, \quad m_{3,17} = m_{17,3} = \frac{\mu_{13}}{2}, \\
m_{3,18} &= m_{18,3} = -\frac{\lambda_{23}}{3}u, \quad m_{3,20} = m_{20,3} = -\frac{\lambda_{23}}{3}w, \\
m_{4,12} &= m_{12,4} = \frac{\lambda_{31}}{3}u, \quad m_{4,15} = m_{15,4} = \frac{\mu_{01}}{2}, \quad m_{4,16} = m_{16,4} = \frac{\lambda_{21}}{3}u, \\
m_{4,19} &= m_{19,4} = \frac{\mu_{11}}{2}, \quad m_{4,20} = m_{20,4} = \frac{\lambda_{31}}{3}v, \\
m_{5,12} &= m_{12,5} = \frac{\lambda_{32}}{3}u, \quad m_{5,15} = m_{15,5} = \frac{\mu_{02}}{2}, \quad m_{5,16} = m_{16,5} = \frac{\lambda_{22}}{3}u, \\
m_{5,19} &= m_{19,5} = \frac{\mu_{12}}{2}, \quad m_{5,20} = m_{20,5} = \frac{\lambda_{32}}{3}v, \\
m_{6,12} &= m_{12,6} = \frac{\lambda_{33}}{3}u, \quad m_{6,15} = m_{15,6} = \frac{\mu_{03}}{2}, \quad m_{6,16} = m_{16,6} = \frac{\lambda_{23}}{3}u, \\
m_{6,19} &= m_{19,6} = \frac{\mu_{13}}{2}, \quad m_{6,20} = m_{20,6} = \frac{\lambda_{33}}{3}v, \\
m_{7,7} &= -m_\lambda, \quad m_{7,12} = m_{12,7} = \frac{gv}{\sqrt{2}}, \quad m_{7,13} = m_{13,7} = -\frac{gv'}{\sqrt{2}}, \\
m_{7,20} &= m_{20,7} = -\frac{gu}{\sqrt{2}}, \quad m_{7,21} = m_{13,7} = -\frac{gu'}{\sqrt{2}}, \\
m_{8,9} &= m_{9,8} = m_{10,10} = -m_\lambda, \quad m_{8,14} = m_{14,8} = gv, \quad m_{8,17} = m_{17,8} = -gw', \\
m_{9,15} &= m_{15,9} = -gv', \quad m_{9,16} = m_{16,9} = gw, \\
m_{10,12} &= m_{12,10} = \frac{gv}{\sqrt{6}}, \quad m_{10,13} = m_{13,10} = -\frac{gv'}{\sqrt{6}}, \\
m_{10,18} &= m_{18,10} = -\sqrt{\frac{2}{3}}gw, \quad m_{10,19} = m_{19,10} = \sqrt{\frac{2}{3}}gw', \\
m_{10,20} &= m_{20,10} = \frac{gu}{\sqrt{6}}, \quad m_{10,21} = m_{21,10} = -\frac{gu'}{\sqrt{6}}, \\
m_{11,11} &= -m', \quad m_{11,12} = m_{12,11} = -\frac{g'v}{3\sqrt{2}}, \quad m_{11,13} = m_{13,11} = \frac{g'v'}{3\sqrt{2}}, \\
m_{11,18} &= m_{18,11} = -\frac{g'w}{3\sqrt{2}}, \quad m_{11,19} = m_{19,11} = \frac{g'w'}{3\sqrt{2}}, \\
m_{11,20} &= m_{20,11} = \sqrt{\frac{2}{3}}g'u, \quad m_{11,21} = m_{21,11} = -\sqrt{\frac{2}{3}}g'u', \\
m_{12,13} &= m_{13,12} = \frac{\mu_\eta}{2}, \quad m_{12,17} = m_{17,12} = \frac{\mu_2}{2}, \\
m_{12,18} &= m_{18,12} = \frac{f_1}{3}u, \quad m_{12,20} = m_{20,12} = \frac{f_1}{3}w, \\
m_{13,16} &= m_{16,13} = \frac{\mu_3}{2}, \quad m_{13,19} = m_{19,13} = \frac{f'_1}{3}u', \quad m_{13,21} = m_{21,13} = \frac{f'_1}{3}w',
\end{aligned}$$

$$\begin{aligned}
m_{14,15} &= m_{15,14} = \frac{\mu_\eta}{2}, \quad m_{14,16} = m_{16,14} = -\frac{f_1}{3}u, \quad m_{14,19} = m_{19,14} = \frac{\mu_2}{2}, \\
m_{15,17} &= m_{17,15} = -\frac{f'_1}{3}u', \quad m_{15,18} = m_{18,15} = \frac{\mu_3}{2}, \\
m_{16,17} &= m_{17,16} = \frac{\mu_\chi}{2}, \\
m_{17,19} &= m_{19,17} = \frac{\mu_\chi}{2}, \quad m_{17,20} = m_{20,17} = \frac{f_1}{3}u, \\
m_{18,19} &= m_{19,18} = \frac{\mu_\chi}{2}, \quad m_{18,21} = m_{21,18} = \frac{f_1}{3}v, \\
m_{19,21} &= m_{21,19} = \frac{f'_1}{3}u', \\
m_{20,21} &= m_{21,20} = \frac{\mu_\rho}{2}.
\end{aligned} \tag{A.1}$$

References

- [1] Y. Fukuda *et al.* (Super-Kamiokande Collaboration), Phys. Rev. Lett. **81**, 1562 (1998); **82**, 2644 (1999); **85**, 3999 (2000); Y. Suzuki, Nucl. Phys. B, Proc. Suppl. **77**, 35 (1999); Y. Fukuda *et al.* (Super-Kamiokande Collaboration), Phys. Rev. Lett. **81**, 1158 (1998); **82**, 1810 (1999); S. Fukuda *et al.*, Phys. Rev. Lett. **86**, 5651 (2001).
- [2] H. Ts. Wong, in *the Proceedings of Hanoi Forum on Frontiers of Basic Science: Towards New Physics: Earth and Space Science, Mathematics*, Hanoi, Vietnam, 27-29 Sep 2005, Osaka University Press (2006), pp. 74 - 79, [arXiv:hep-ex/0510001].
- [3] G. L. Fogli, E. Lisi, A. Marrone and A. Palazzo, [arXiv:hep-ph/0506083].
- [4] M. Gell-Mann, P. Ramond and R. Slansky, in *Super-gravity, Proceedings of the Workshop*, Stony Brook, New York, 1979, edited by D. Freedman and P. van Nieuwenhuizen (North-Holland, Amsterdam, 1979); J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980); **25**, 774 (1982).
- [5] A. Zee, Phys. Lett. B **93**, 389 (1980); Nucl. Phys. B **264**, 99 (1986); K. S. Babu, Phys. Lett. B **203**, 132 (1988); K. S. Babu and E. Ma, Phys. Rev. Lett. **61**, 674 (1988); D. Chang, W.-Y. Keung and P. B. Pal, Phys. Rev. Lett. **61**, 2420 (1988).
- [6] F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992); P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992); R. Foot *et al.*, Phys. Rev. D **47**, 4158 (1993).
- [7] M. Singer, J. W. F. Valle and J. Schechter, Phys. Rev. D **22**, 738 (1980); R. Foot, H. N. Long and Tuan A. Tran, Phys. Rev. D **50**, 34(R) (1994); J. C. Montero, F. Pisano and V. Pleitez, Phys. Rev. D **47**, 2918 (1993); H. N. Long, Phys. Rev. D **53**, 437 (1996); Phys. Rev. D **54**, 4691 (1996).

- [8] F. Pisano, Mod. Phys. Lett. A **11**, 2639 (1996); C. A. de S. Pires and O. P. Ravinez, Phys. Rev. D **58**, 035008 (1998); J. C. Montero, C. A. de S. Pires and V. Pleitez, Phys. Lett. B **502**, 167 (2001); A. G. Dias and V. Pleitez, Phys. Rev. D **69**, 077702 (2004); A. G. Dias and A. Doff, Phys. Rev. D **72**, 035006 (2005).
- [9] T. Kitabayashi and M. Yasuè, Phys. Rev. D **63**, 095002 (2001); Phys. Lett. B **508**, 85 (2001); Phys. Rev. D **63**, 095006 (2001); Y. Okamoto and M. Yasuè, Phys. Lett. B **466**, 267 (1999).
- [10] H. N. Long and T. Inami, Phys. Rev. D **61**, 075002 (2000); P. V. Dong, H. N. Long, D. T. Nhung and D. V. Soa, Phys. Rev. D **73**, 035004 (2006); P. V. Dong, H. N. Long, and D. V. Soa, *Higgs-gauge boson interactions in the economical 3-3-1 model*, to be published in Phys. Rev. D, [arXiv:hep-ph/0603108].
- [11] D. Chang and H. N. Long, Phys. Rev. D **73**, 053006 (2006).
- [12] A. G. Dias, C. A. de S. Pires and P. S. Rodrigues da Silva, Phys. Lett. B **628**, 85 (2005).
- [13] J. C. Montero, V. Pleitez and M. C. Rodriguez, Phys. Rev. D **70**, 075004 (2004); D. T. Huong, M. C. Rodriguez and H. N. Long, [arXiv:hep-ph/0508045].
- [14] J. C. Montero, V. Pleitez and M. C. Rodriguez, Phys. Rev. D **65**, 035006 (2002).
- [15] J. C. Montero, V. Pleitez and M. C. Rodriguez, Phys. Rev. D **65**, 095008 (2002).
- [16] A. Salam and J. Strathdee, Nucl. Phys. B **87**, 85 (1975).
- [17] P. Fayet, Nucl. Phys. B **90**, 104 (1975).
- [18] R. Barbier *et al.*, Phys. Rept. **420**, 1 (2005), [arXiv:hep-ph/0406039].
- [19] G. Moreau, [arXiv:hep-ph/0012156].
- [20] D. Bailin and A. Love, *Supersymmetric Gauge Field Theory and String Theory*, 1st edition, Institute of Physics Publishing, Bristol UK, 1994.
- [21] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, 2nd edition, Princeton University Press, Princeton NJ, 1992.
- [22] H. E. Haber and G. L. Kane, Phys. Rept. **117**, 75 (1985).
- [23] L. Girardello and M. T. Grisaru, Nucl. Phys. B **194**, 65 (1982).
- [24] P. Fayet, *New Frontiers in High-Energy Physics*, Proc. Orbis Scientiae, Coral Gables (Florida, USA), 1978, edited by A. Perlmutter and L.F. Scott (Plenum, N.Y., 1978), pp. 413.
- [25] G. R. Farrar and P. Fayet, Phys. Lett. B **76**, 575 (1978).

- [26] L. Hall and M. Suzuki, Nucl. Phys. B **231**, 419 (1984).
- [27] L. E. Ibañez and G. G. Ross, Nucl. Phys. B **368**, 3 (1992).
- [28] L. J. Hall and M. Suzuki, Nucl. Phys. B **231**, 419 (1984).
- [29] T. Banks, Y. Grossman, E. Nardi and Y. Nir, Phys. Rev. D **52**, 5319 (1995).
- [30] M. A. Diaz, J. C. Romão and J. W. F. Valle, Nucl. Phys. B **524**, 23 (1998).
- [31] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).
- [32] M. Capdequi-Peyranère and M. C. Rodriguez, Phys. Rev. D **65**, 035006 (2002).
- [33] M. Drees, R. Godbole and P. Roy, *Theory and Phenomenology of Sparticles*, World Scientific Publishing, Singapore, 2004.
- [34] L. Roszkowski, [arXiv:hep-ph/9302259].